



## King's Research Portal

DOI:

[10.1016/j.jacbts.2019.01.002](https://doi.org/10.1016/j.jacbts.2019.01.002)

*Document Version*

Publisher's PDF, also known as Version of record

[Link to publication record in King's Research Portal](#)

*Citation for published version (APA):*

Ueda, H., Yamaguchi, O., Taneike, M., Akazawa, Y., Wada-Kobayashi, H., Sugihara, R., Yorifuji, H., Nakayama, H., Omiya, S., Murakawa, T., Sakata, Y., & Otsu, K. (2019). Administration of a TLR9 inhibitor attenuates the development and progression of heart failure in mice. *JACC: Basic to Translational Science*, 4(3), 348-363. [10]. <https://doi.org/10.1016/j.jacbts.2019.01.002>

### **Citing this paper**

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

### **General rights**

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

### **Take down policy**


If you believe that this document breaches copyright please contact [librarypure@kcl.ac.uk](mailto:librarypure@kcl.ac.uk) providing details, and we will remove access to the work immediately and investigate your claim.

## PRECLINICAL RESEARCH

# Administration of a TLR9 Inhibitor Attenuates the Development and Progression of Heart Failure in Mice



Hiromichi Ueda, MD,<sup>a</sup> Osamu Yamaguchi, MD, PhD,<sup>a</sup> Manabu Taneike, MD, PhD,<sup>a</sup> Yasuhiro Akazawa, MD,<sup>a</sup> Haruko Wada-Kobayashi, MD,<sup>a</sup> Ryuta Sugihara, MD,<sup>a</sup> Hiroki Yorifuji, MD,<sup>a</sup> Hiroyuki Nakayama, MD, PhD,<sup>b</sup> Shigemiki Omiya, MD, PhD,<sup>c</sup> Tomokazu Murakawa, MD, PhD,<sup>c</sup> Yasushi Sakata, MD, PhD,<sup>a</sup> Kinya Otsu, MD, PhD<sup>c</sup>



## HIGHLIGHTS

- Under pressure overload, mitochondrial deoxyribonucleic acid containing the unmethylated cytidine-phosphate-guanosine motif is accumulated in cardiomyocytes and stimulates Toll-like receptor 9, resulting in inflammation and heart failure.
- Treatment with E6446, (6-[3-(pyrrolidin-1-yl)propoxy]-2-(4-(3-(pyrrolidin-1-yl)propoxy)phenyl]benzo[d]oxazole), a specific Toll-like receptor 9 inhibitor, prevented the development and slowed the progression of left ventricular dilatation and cardiac dysfunction in mice after pressure overload.
- E6446 attenuated the inflammatory responses in the pressure-overloaded mouse heart, even though the accumulation of mitochondrial deoxyribonucleic acid in cardiomyocytes was observed.
- E6446 could be a new therapeutic agent against heart failure.

From the <sup>a</sup>Department of Cardiovascular Medicine, Graduate School of Medicine, Osaka University, Suita, Osaka, Japan;

<sup>b</sup>Laboratory of Clinical Science and Biomedicine, Graduate School of Pharmaceutical Sciences, Osaka University, Suita, Osaka, Japan; and the <sup>c</sup>School of Cardiovascular Medicine and Sciences, King's College London British Heart Foundation Centre of Excellence, London, United Kingdom. This research was supported by the British Heart Foundation (CH/11/3/29051 and RG/16/15/32294), Fondation Leducq (RA15CVD04), and JSPS KAKENHI (18H02807), to Dr. Otsu. The authors have reported that they have no relationships relevant to the contents of this paper to disclose.

All authors attest they are in compliance with human studies committees and animal welfare regulations of the authors' institutions and Food and Drug Administration guidelines, including patient consent where appropriate. For more information, visit the JACC: Basic to Translational Science [author instructions page](#).

Manuscript received October 3, 2018; revised manuscript received January 10, 2019, accepted January 11, 2019.

## SUMMARY

Mitochondrial deoxyribonucleic acid, containing the unmethylated cytidine-phosphate-guanosine motif, stimulates Toll-like receptor 9 to induce inflammation and heart failure. A small chemical, E6446 [(6-[3-(pyrrolidin-1-yl)propoxy]-2-(4-(3-(pyrrolidin-1-yl)propoxy)phenyl]benzo[d]oxazole)], is a specific Toll-like receptor 9 inhibitor in cardiomyocytes. In this study, we showed that E6446 exerts beneficial effects for the prevention and treatment of pressure overload-induced heart failure in mice. When administered before the operation and chronically thereafter, E6446 prevented the development of left ventricular dilatation as well as cardiac dysfunction, fibrosis, and inflammation. Furthermore, when administered after the manifestation of cardiac dysfunction, E6446 slowed progression of cardiac remodeling. Thus, the inhibitor may be a novel therapeutic agent for treating patients with heart failure. (J Am Coll Cardiol Basic Trans Science 2019;4:348-63) © 2019 The Authors. Published by Elsevier on behalf of the American College of Cardiology Foundation. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

Heart failure is a complex disease associated with high levels of morbidity and mortality and marked reductions in quality of life. Previous extensive studies on heart failure have reported an important role for proinflammatory cytokines in its pathogenesis (1). Circulating levels of proinflammatory cytokines, including tumor necrosis factor (TNF)- $\alpha$ , are related to the severity and prognosis of the disease. However, the targeted anti-TNF- $\alpha$  approaches were neutral with respect to the primary endpoints of the trial or resulted in worsening heart failure or death (2,3). In addition to TNF- $\alpha$ , the pro-inflammatory cytokines that are elaborated in heart failure include other members of the TNF superfamily, members of the interleukin-1 family, and interleukin (IL)-6 (1). Recognizing the molecular mechanism underlying the developing inflammation in heart failure is essential for developing strategies to control disease progression, including therapeutic drugs.

Mitochondrial deoxyribonucleic acid (DNA) contains the unmethylated cytidine-phosphate-guanosine (CpG) motif, which stimulates Toll-like receptor (TLR) 9 to induce inflammation (4,5). Mitochondria damaged by external hemodynamic stress are degraded by the autophagy/lysosome system in cardiomyocytes (6). Insufficient degradation of mitochondrial DNA mediated through autophagy in pressure-overloaded mouse hearts leads to its binding to TLR9 to induce inflammation and heart failure (7). In failing mouse hearts, mitochondrial DNA is located in autolysosomes. Furthermore, TLR9 ablation in pressure-overloaded mice attenuated the development of inflammation and heart failure. Thus, interference with TLR9 function by small molecules is likely to produce a better clinical outcome by preventing its aberrant inflammatory responses.

E6446 (6-[3-(pyrrolidin-1-yl)propoxy]-2-(4-(3-(pyrrolidin-1-yl)propoxy)phenyl]benzo[d]oxazole), is a synthetic antagonist of nucleic acid-sensing TLRs and is orally bioactive (8,9). In vitro, E6446 specifically inhibits the activation of TLR9 (8). Others have reported that the compound inhibits TLR9 but also TLR7 in a ligand-dependent manner (9). When E6446 is administered to mice, it suppresses inflammatory responses to challenge doses of unmethylated CpG containing oligodeoxynucleotide (CpG ODN) (8,9). When E6446 is administered chronically in mouse cerebral malaria and spontaneous lupus models, the compound inhibits cytokine production with prevention of signs of cerebral malaria and circulating antinuclear antibodies, respectively.

In the present study, the efficacy of oral treatment with E6446 was evaluated on mouse pressure overload-induced heart failure models. Our results indicate that E6446 exerts beneficial effects for the prevention and treatment of heart failure in mice and might be a novel therapeutic agent for treating patients with heart failure.

## METHODS

**CELL CULTURE.** Adult cardiomyocytes were isolated from 10- to 12-week-old C57BL/6J male mice (CLEA Japan, Inc., Tokyo, Japan) by using a Langendorff system and were then cultured (7).

**RIBONUCLEIC ACID ANALYSIS.** Total ribonucleic acid (RNA) was extracted from the left ventricle or cultured cardiomyocytes by using the TRIzol reagent (Thermo Fisher Scientific, Waltham, Massachusetts) and reverse transcribed by using TaqMan Reverse Transcription Reagents (Thermo Fisher Scientific) (7).

## ABBREVIATIONS AND ACRONYMS

- CCCP** = carbonyl cyanide *m*-chlorophenyl hydrazine
- CpG** = cytidine-phosphate-guanosine
- CpG ODN** = unmethylated cytidine-phosphate-guanosine containing oligodeoxynucleotide
- DNA** = deoxyribonucleic acid
- E6446** = (6-[3-(pyrrolidin-1-yl)propoxy]-2-(4-(3-(pyrrolidin-1-yl)propoxy)phenyl]benzo[d]oxazole)
- EdU** = 5-ethynyl-2'-deoxyuridine
- IL** = interleukin
- IVsd** = end-diastolic interventricular septal wall thickness
- LAMP** = lysosome-associated membrane protein
- LC** = microtubule-associated protein light chain
- LPS** = lipopolysaccharide
- LV** = left ventricular
- mRNA** = messenger ribonucleic acid
- TAC** = transverse aortic constriction
- TLR** = Toll-like receptor
- TNF** = tumor necrosis factor

**TABLE 1 Echocardiographic Parameters of All Mice Subjected to TAC Operation for 2 Weeks**

	Baseline (n = 30)	TAC for 2 Weeks (n = 30)
LVDd, mm	2.31 ± 0.02	2.77 ± 0.03*
LVDs, mm	0.88 ± 0.01	1.45 ± 0.03*
LVFS, %	62.0 ± 0.40	47.8 ± 0.71*
IVSd, mm	0.92 ± 0.00	1.04 ± 0.01*
LVPWd, mm	0.86 ± 0.01	1.02 ± 0.01*
Heart rate, beats/min	696 ± 3.4	685 ± 4.3
LV mass, mg	58.7 ± 0.7	96.6 ± 2.1*

The echocardiographic parameters of the mice were obtained 2 weeks after the operation. Values are mean ± SE. \*p < 0.05 vs. baseline.

IVSd = end-diastolic interventricular septal wall thickness; LV = left ventricular; LVDd = end-diastolic left ventricular internal dimension; LVDs = end-systolic left ventricular internal dimension; LVFS = left ventricular fractional shortening; LVPWd = end-diastolic left ventricular posterior wall thickness; TAC = transverse aortic constriction.

Real-time quantitative polymerase chain reaction was performed by using the Platinum Quantitative PCR SuperMix-UDG (Thermo Fisher Scientific). Relative levels of gene expression were normalized to the *Gapdh* messenger RNA (mRNA) expression. The primers (Thermo Fisher Scientific: Assay identity) used were as follows: *Nppa*, Mm01255747\_g1; *Nppb*, Mm00435304\_g1; *Col1a2*, Mm01165107\_m1; *Col3a1*, Mm00802331\_m1; *Gapdh*, 4352339E; *Il6*, Mm99999064\_m1; *Il1b*, Mm01336189\_m1; and *Tnfa*, Mm00443260\_g1.

**IMMUNOFLUORESCENCE MICROSCOPY.** Adult mouse cardiomyocytes on laminin-coated glass-based dishes (IWAKI Cell Biology, Bio-REV Pte. Ltd., Singapore) were incubated with 100 nmol/l carbonyl cyanide *m*-chlorophenyl hydrazine (CCCP) for 6 h. To estimate mitochondrial membrane potential, the cells were treated with 10 nmol/l of tetramethylrhodamine ethyl ester (Molecular Probes, Eugene, Oregon) for 30 min. To visualize DNA and autophagosomes, the cells were incubated in three-dimensional gel with Cellmatrix I-A (Nitta Gelatin Inc., Osaka, Japan) and fixed with methanol at −30°C for 15 min. The cells were incubated with anti-microtubule-associated protein light chain (LC) 3B antibody (Cell Signaling Technology, Danvers, Massachusetts) overnight at 4°C, followed by staining with anti-rabbit Alexa 568 secondary antibody (Abcam, Cambridge, United Kingdom) overnight. The cells were incubated with 100-fold diluted PicoGreen (Thermo Fisher Scientific) for 30 min before confocal microscopic analysis using an FV-1000D microscope (Olympus, Tokyo, Japan) (7).

**ANIMAL STUDY.** The investigation conforms to the Position of the American Heart Association on Research Animal Use adopted by the American Heart Association on November 11, 1984. All in vivo and


in vitro experimental protocols were conducted under the supervision of the Animal Research Committee of Osaka University and in accordance with the Guidelines for Animal Experiments of Osaka University and the Japanese Animal Protection and Management Law.

The 10- to 12-week-old male C57BL/6J mice were subjected to transverse aortic constriction (TAC) surgery (10). Sham-operated animals underwent the same operation without aortic constriction. Blood pressure was measured noninvasively on mice anesthetized with 2.5% tribromoethanol by using a pressure monitor (Model MK-2000, Muromachi Kikai Co., Ltd., Tokyo, Japan). The pressure gradient across TAC was estimated by the difference in blood pressure between both arms by using a pressure monitor. Ultrasonography (Sonos 5500, equipped with a 15-MHz linear probe, Philips Medical Systems, Cambridge, Massachusetts) was used for assessing left ventricular (LV) size and function on conscious mice.

**DRUG AND TREATMENT.** E6446 (Eisai, Inc., Andover, Massachusetts) was dissolved in dimethyl sulfoxide (final 0.04% v/v) for experiments. Isolated cardiomyocytes were pretreated with E6446 (0 to 10 μmol/l) for 1 h, followed by treatment with 1 μg/ml of lipopolysaccharide (LPS) (FUJIFILM WAKO Pure Chemical Co., Osaka, Japan), 2 mmol/l of loxoribine (InvivoGen, San Diego, California), or 5 μmol/l of type B CpG ODN (ODN1668, InvivoGen) for 6 h. To examine the effect of E6446 on the level of cytokine mRNAs, cardiomyocytes were pretreated with 10 μmol/l of E6446 for 1 h, followed by treatment with 100 nmol/l of CCCP for 6 h.

For the in vivo ODN1668 challenge experiments with E6446, the mice were orally administered E6446 at a dose of 1.5 mg/200 μl per mouse using animal feeding needles (Natsume Seisakusho Co., Ltd., Tokyo, Japan). One, two, or 3 days later, the mice were injected intraperitoneally with 60 μg/mouse of ODN1668 2 h before sacrifice, following intraperitoneal injection of 20 mg/mouse of D-galactosamine (MilliporeSigma, Burlington, Massachusetts) for 30 min. In the prevention study, E6446 administration was started 2 days before the operation. Mice were administered E6446 orally at a dose of 1.5 mg/mouse or its vehicle (saline) as a control every 2 days. In the treatment study, 30 mice were subjected to TAC operation for 2 weeks. Ten mice with fractional shortening >50% were excluded from the study. The remaining 20 mice were randomized to the saline- and E6446-treated groups and then orally administered saline or E6446 (1.5 mg/mouse) every 2 days (Table 1).

**FIGURE 1** Selectivity of E6446




The expression level of cytokine messenger ribonucleic acid in cardiomyocytes with increasing concentrations of E6446, (6-[3-(pyrrolidin-1-yl)propoxy]-2-(4-(3-(pyrrolidin-1-yl)propoxy)phenyl]benzo[d]oxazole) ( $n = 3$ ). Cardiomyocytes were treated with 0 to 10  $\mu\text{M}$ /l of E6446 for 1 h, followed by (A) 5  $\mu\text{M}$ /l ODN1668, (B) 1  $\mu\text{g}/\text{ml}$  of lipopolysaccharide (LPS), or (C) 2  $\text{mmol}/\text{l}$  of loxoribine for 6 h. Control groups were treated with vehicle. Data were normalized to the content of *Gapdh* messenger ribonucleic acid. Values are mean  $\pm$  SE. Bars in graphs indicate  $p < 0.05$ . \* $p < 0.05$  versus both control groups.

A blood sample was taken from the right ventricle. IL-6, IL-1B, and TNF- $\alpha$  serum levels were measured with an enzyme-linked immunoadsorbent assay kit (Thermo Fisher Scientific for IL-6 and IL-1B, R and D Systems [Minneapolis, Minnesota] for TNF- $\alpha$ ).

**HISTOLOGICAL ANALYSIS.** Heart samples were fixed in buffered 4% paraformaldehyde solution and embedded in paraffin (10). Fibrosis fraction was quantified by using ImageJ software (National Institutes of Health, Bethesda, Maryland) and


**FIGURE 2** Effect of E6446 on Mitochondrial Damage-Induced Cytokine Messenger Ribonucleic Acid Production in Isolated Adult Mouse Cardiomyocytes




(A) Disruption of mitochondrial membrane potential by carbonyl cyanide *m*-chlorophenyl hydrazine (CCCP). Isolated cardiomyocytes were incubated with 100 nmol/L CCCP for 6 h and stained with tetramethylrhodamine ethyl ester (TMRE). Scale bar, 20  $\mu$ m. The right graph shows the proportion of TMRE-positive cells. (B) Double staining of E6446-treated cardiomyocytes stimulated with CCCP using PicoGreen (green) and anti-microtubule-associated protein light chain (LC) 3B antibody (red). Low-magnified images are shown in the left panels. Scale bar, 20  $\mu$ m. Higher magnified images of the squared areas are shown in the right panels. Scale bar, 5  $\mu$ m. Arrows indicate PicoGreen and LC3B merged deposits. The right graph shows the number of PicoGreen and LC3B double-positive deposits per cell. Ten or more cells were analyzed for each experiment. (C) Inhibition of cytokine messenger ribonucleic acid expression in CCCP-stimulated cardiomyocytes by E6446. Cardiomyocytes were treated with 10  $\mu$ mol/L E6446 for 1 h, followed by 100 nmol/L CCCP for 6 h. Values are mean  $\pm$  SE (n = 3). \*p < 0.05 versus corresponding controls. †p < 0.05 versus all other groups. Abbreviation as in Figure 1.



**FIGURE 3** Determination of the Experimental Conditions for E6446 Administration





The time dependence of (A) cytokine protein and (B) messenger ribonucleic acid (mRNA) expression in the heart after injection with ODN1668. Mice (body weight 24.4 to 25.8 g) were pretreated with oral administration of 1.5 mg/mouse (60 mg/kg) of E6446 1, 2, or 3 days before intraperitoneal injection with 60  $\mu$ g/mouse of ODN1668. Two hours after ODN1668 injection, mice were sacrificed for analysis (see Figure 9A). Dose dependency in the inhibition of (C) cytokine protein and (D) mRNA expression after ODN1668 injection with increasing concentrations of E6446. Two days after oral administration with the indicated dose of E6446, mice were administered an intraperitoneal injection of 60  $\mu$ g/mouse of ODN1668. Two hours later, mice were sacrificed for analysis (see Figure 9B). Data were normalized to the content of *Gapdh* mRNA in B and D. Values are mean  $\pm$  SE (n = 3). Bars in graphs indicate p < 0.05. IL = interleukin; TNF = tumor necrosis factor; other abbreviation as in Figure 1.

**FIGURE 4** Improvement of Echocardiographic Parameters by Treatment With E6446 Initiated Before TAC

The oral administration of E6446 (1.5 mg/mouse) was performed every 2 days from 2 days before transverse aortic constriction (TAC) (see [Figure 10A](#)). **(A)** Pressure gradient across TAC estimated by using a pressure monitor 1 week after operation. **(B)** Representative images of transthoracic M-mode echocardiographic tracing. Scale bars, 0.2 s and 5 mm, respectively. **(C)** Echocardiographic parameters of the mice treated with E6446 4 weeks after TAC (n = 4 in sham groups, n = 5 in TAC groups). Values are mean  $\pm$  SE. \*p < 0.05 versus sham-operated groups. †p < 0.05 versus all other groups. FC = fractional shortening; IVSd = end-diastolic interventricular septal wall thickness; LV = left ventricular; LVDd = end-diastolic left ventricular internal dimension; LVDs = end-systolic left ventricular internal dimension; LVPWd = end-diastolic left ventricular posterior wall thickness; other abbreviation as in [Figure 1](#).







**FIGURE 6** Effects of E6446 on mRNA Expressions

The treatment with E6446 was performed every 2 days from 2 days before TAC. The mice were analyzed 4 weeks after the operation (see Figure 10A). Data were normalized to the content of *Gapdh* mRNA. Values are mean  $\pm$  SE ( $n = 4$  in sham groups,  $n = 5$  in TAC groups). \* $p < 0.05$  versus all other groups. Abbreviations as in Figures 1, 3, and 4.

expressed as a proportion of the ventricles. For immunohistochemical analysis, frozen heart sections 5- $\mu$ m thick were fixed in buffered paraformaldehyde and stained with anti-mouse CD45 (R and D Systems), CD68 (Bio-Rad, Hercules, California), Ly6G/C (BD Pharmingen, BD Biosciences, San Jose, California), CD3 (Abcam), and lysosome-associated membrane

protein (LAMP) 2a (Thermo Fisher Scientific) antibodies. To measure the cardiomyocyte cross-sectional area, the frozen heart sections were stained with WGA-Alexa 555 (Thermo Fisher Scientific) for 1 h. The cardiomyocyte cross-sectional area was measured by tracing the outline of >100 cardiomyocytes in each section by using ImageJ software. For detection of



**FIGURE 8** Improvement of Echocardiographic Parameters and Lung Congestion After Treatment With E6446 Initiated After TAC

Two weeks after TAC, treatment with E6446 was performed every 2 days (see [Figure 10B](#)). Values are mean  $\pm$  SE ( $n = 10$ ). **(A)** Pressure gradient across TAC estimated by using a pressure monitor 1 week after operation. **(B)** Representative images of transthoracic M-mode echocardiographic tracing. Scale bars, 0.2 s and 5 mm, respectively. **(C)** Echocardiographic parameters. The parameters were examined for 6 weeks after TAC. **Open and closed circles** indicate saline-treated control groups and E6446-treated groups, respectively. The data were analyzed by using 2-way repeated measure analysis of variance followed by Tukey's post hoc test. \* $p < 0.05$  between the 2 groups at the corresponding time point. **Dotted lines** indicate  $p < 0.05$  between the 2 saline-treated control groups at different time points. **Solid lines** indicate  $p < 0.05$  between the 2 E6446-treated groups at different time points. **(D)** Physiological parameters 6 weeks after the operation. \* $p < 0.05$  versus saline-treated control group. Abbreviations as in [Figures 1, 4, and 5](#).

mitochondrial DNA, mice were intraperitoneally injected with 5 mg of 5-ethynyl-2'-deoxyuridine (EdU) (Invitrogen) 1 day before sacrifice. EdU was detected by using a Click-iT EdU Alexa Fluor 488 Imaging Kit (Thermo Fisher Scientific) (7). EdU- and LAMP2a double-positive deposits were counted in 5 different areas per section in each mouse.

**STATISTICAL ANALYSIS.** Results are shown as mean  $\pm$  SE. GraphPad Prism version 7.04 (GraphPad Software, La Jolla, California) was used for statistical analysis. A Student's *t*-test was used for a 2-group comparison; a 1-way analysis of variance followed by Tukey's post hoc test or 2-way repeated measure analysis of variance followed by Tukey's post hoc test were used for multiple comparisons. Significant differences were defined as  $p < 0.05$ .

## RESULTS

**EFFECT OF E6446 ON CYTOKINE mRNA PRODUCTION IN ISOLATED CARDIOMYOCYTES.** Isolated cardiomyocytes were stimulated by a TLR9 ligand (e.g., ODN1668), a TLR4 ligand (e.g., LPS), or a TLR7 ligand (e.g., loxoribine) in the presence of increasing concentrations of E6446 (11,12). ODN1668 significantly increased the expression levels of *Il6*, *Il-1B* (*Il1b*), and *Tnfa* mRNAs (Figure 1A). Incubation of cardiomyocytes with E6446 significantly reduced the induction of *Il6*, *Il1b*, and *Tnfa* mRNAs in response to ODN1668. LPS significantly increased the expression levels of *Il6*, *Il1b*, and *Tnfa* mRNAs, whereas loxoribine significantly increased the expression levels of *Il1b* and *Tnfa* mRNAs but not *Il6* mRNA (Figures 1B and 1C). E6446 had no effect on the induction of the cytokine mRNAs induced by LPS or loxoribine.

Incubation of cardiomyocytes with CCCP diminished mitochondrial membrane potential (Figure 2A). Cardiomyocytes were stained with PicoGreen, a highly sensitive marker for DNA, and anti-LC3B antibody, a marker for autophagosomes. CCCP increased the number of PicoGreen and LC3B double-positive deposits, suggesting accumulation of DNA in autophagosomes or autolysosomes (Figure 2B). Incubation of cardiomyocytes with CCCP increased the levels of *Il6* and *Il1b* mRNAs (Figure 2C). E6446 had no effect on the number of PicoGreen and LC3B double-positive deposits, but it significantly reduced *Il6* and *Il1b* mRNA expression in CCCP-treated cardiomyocytes.

**IN VIVO ADMINISTRATION OF E6446.** To examine the in vivo efficacy of E6446 on inhibition of TLR9 and determine the experimental conditions for its administration in mice, 1.5 mg/mouse (60 mg/kg) of E6446 was orally administered to mice 3, 2, or 1 day before intraperitoneal injection of ODN1668. Two

**TABLE 2** Echocardiographic Parameters of Mice Included in the E6446 Treatment Study 2 Weeks After TAC Operation

	Baseline		2 Weeks After TAC	
	Saline (n = 10)	E6446 (n = 10)	Saline (n = 10)	E6446 (n = 10)
LVDd, mm	2.30 $\pm$ 0.03	2.27 $\pm$ 0.03	2.81 $\pm$ 0.03*	2.86 $\pm$ 0.03*
LVDs, mm	0.88 $\pm$ 0.02	0.86 $\pm$ 0.02	1.54 $\pm$ 0.03*	1.55 $\pm$ 0.02*
LVFS, %	61.5 $\pm$ 0.71	62.3 $\pm$ 0.79	45.2 $\pm$ 0.77*	45.6 $\pm$ 0.43*
IVSd, mm	0.92 $\pm$ 0.01	0.92 $\pm$ 0.01	1.07 $\pm$ 0.01*	1.05 $\pm$ 0.01*
LVPWd, mm	0.86 $\pm$ 0.02	0.87 $\pm$ 0.01	1.04 $\pm$ 0.01*	1.03 $\pm$ 0.01*
Heart rate, beats/min	692 $\pm$ 3.9	695 $\pm$ 6.5	689 $\pm$ 7.2	672 $\pm$ 7.5
LV mass, mg	58.2 $\pm$ 1.4	58.9 $\pm$ 1.1	103.5 $\pm$ 3.1*	101.4 $\pm$ 2.1*

Values are mean  $\pm$  SE. Thirty mice were subjected to TAC operation for 2 weeks. Ten mice with fractional shortening  $>50\%$  were excluded from the study. The remaining 20 mice were randomized to the saline- and E6446-treated groups. The parameters of the mice were obtained 2 weeks after the operation by using echocardiography. \* $p < 0.05$  vs. corresponding control at baseline.


Abbreviations as in Table 1.

hours after the mice were administered ODN, the treated mice produced a higher level of IL-6 than control mice (Figure 3A). When E6446 was administered 2 or 1 day before ODN1668 injection, IL-6 levels were lower than the E6446-nontreated ODN1668-injected group and showed no significant difference compared with the control group without ODN1668 injection. When E6446 was administered 3 days before ODN1668 injection, the level of IL-6 did not differ from that in the nontreated group. To confirm the protein data, mRNA levels in the heart were measured (Figure 3B). The level of *Il6* mRNA in ODN1668-injected mouse hearts was higher than that in control hearts. When E6446 was administered 2 or 1 day before ODN1668 injection, *Il6* mRNA levels were lower than the E6446-nontreated ODN1668-injected group. When E6446 was administered 3 days before ODN1668 injection, the level of *Il6* mRNA differed from the control group but not from other ODN1668-injected groups. The level of *Il1b* mRNA in the E6446-nontreated ODN1668-injected mouse hearts was higher than that in the control hearts and showed a significant difference from that treated with E6446 1 and 2 days but not 3 days before ODN1668 injection; the levels of IL-1B protein were not significantly different among groups. The level of TNF- $\alpha$  in the ODN1668-injected mouse group was higher than that in the nontreated control group and in all the E6446-treated ODN1668-injected groups. The level of *Tnfa* mRNA in E6446-nontreated ODN1668-injected mouse hearts was higher than that in control hearts but showed no difference from all other groups. Thus, the inhibitory effect of E6446 on the induction of cytokines lasted over a period of 2 days, providing a rationale for every-other-day dosing.

The dose of E6446 necessary to inhibit the induction of cytokines was next examined. The serum level



**FIGURE 9 Schematic Protocol to Determine the Experimental Conditions for E6446 Administration**



(A) Schematic protocol to examine the time dependence of cytokine protein and mRNA expression in the heart after injection with ODN1668 (n = 3 for each group). Mice were pretreated with E6446 1.5 mg/mouse for 1, 2, or 4 days before injection of 60 μg/mouse of ODN1668. Two hours after ODN1668 injection, mice were sacrificed for analysis, following treatment with D-galactosamine. (B) Schematic protocol to examine dose dependency in the inhibition of cytokine expression after ODN1668 injection with increasing concentrations of E6446 (n = 3 for each group). Two days after administration with the indicated dose of E6446, mice received 60 μg/mouse of ODN1668, following D-galactosamine injection. Two h later, mice were sacrificed for analysis. Abbreviations as in Figures 1 and 3.

of IL-6 and TNF-α protein in mice treated with 1.5 or 15 mg/mouse of E6446 was lower than that in non-treated mice or in mice treated with 0.15 mg/mouse of E6446 but did not differ from that in control mice (Figure 3C). ODN1668 injection increased serum levels of IL-1B protein. E6446 administration produced no significant inhibitory effect on the increase of IL-1B protein. The levels of *Il6*, *Il1b*, and *Tnfa* mRNAs in hearts treated with 1.5 or 15 mg/mouse were significantly lower than those in the hearts of nontreated mice (Figure 3D). Thus, 1.5 mg/mouse of E6446 was administered every 2 days in the following experiments.

## PREVENTION OF THE DEVELOPMENT OF HEART FAILURE BY E6446

To investigate the efficacy of E6446 on the development of heart failure, mice orally received E6446 or saline 2 days before TAC and every 2 days for 4 weeks thereafter. There was no significant difference in pressure gradient across TAC between the E6446- and saline-treated groups 1 week after TAC (Figure 4A). Four weeks after TAC, saline-treated mice exhibited larger end-diastolic LV internal dimensions and end-systolic LV internal dimensions and lower fractional shortening than those in the sham-operated saline-treated group (Figures 4B and 4C). E6446 treatment significantly reduced LV chamber size and improved cardiac function in TAC-operated mice. TAC increased end-diastolic interventricular septal wall thickness (IVSd) and end-diastolic LV posterior wall thickness in saline- and E6446-treated mice. However, there were no significant differences in IVSd and end-diastolic LV posterior wall thickness between saline- and E6446-treated mice. LV mass was increased by TAC, and E6446 treatment attenuated the increase in LV mass in TAC-operated mice.

There was no significant difference in body weight between the 4 groups (Figure 5A). TAC-operated saline-treated mice exhibited increases in the heart weight-to-tibia length ratio and the lung weight-to-tibia length ratio. E6446 significantly attenuated cardiac hypertrophy and lung congestion. TAC increased the cross-sectional area of cardiomyocytes in saline-treated mice, and the increase was significantly attenuated by E6446 treatment (Figure 5B). *Nppa* and *Nppb* mRNAs increased in TAC-operated saline-treated mice. E6446 attenuated the increases induced by TAC (Figure 6). TAC-operated saline-treated mice exhibited cardiac fibrosis, which was diminished in E6446-treated mice (Figure 5C). The mRNA levels of *Col1a2* and *Col3a1* increased in TAC-operated saline-treated mice. E6446 attenuated the increase in the mRNAs.

## ATTENUATION OF INFLAMMATION IN PRESSURE-OVERLOADED HEARTS BY E6446

TAC-operated saline-treated mice showed infiltration of CD45<sup>+</sup> cells, including CD68<sup>+</sup> macrophages in the heart, which was inhibited by treatment with E6446 (Figure 7A). Although increases in the mRNA expressions of *Il6* and *Tnfa* were detected in saline-treated TAC-operated mice, E6446 attenuated the induction of the mRNAs (Figure 6).

To label mitochondrial DNA, mice were injected with EdU 1 day before sacrifice. EdU specifically binds to mitochondrial DNA during active DNA synthesis in nondividing cardiomyocytes (7). LAMP2a is a marker for lysosomes. In TAC-operated saline- and




E6446-treated hearts, EdU and LAMP2a double-positive deposits were observed (Figure 7B). There was no significant difference in the number of the double-positive deposits between TAC-operated saline- and E6446-treated hearts.

**SLOWING THE PROGRESSION OF HEART FAILURE BY E6446.** Finally, the effect of E6446 on the progression of an already established disease was examined. Mice were subjected to TAC operation (Figure 8A) and divided into 2 groups 2 weeks later. There were no significant differences in echocardiographic parameters between the 2 groups, which already exhibited chamber dilatation and cardiac dysfunction (Table 2). The mice were then administered E6446 or saline every 2 days for 4 weeks. LV chamber dilatation and cardiac dysfunction progressively worsened with time in both groups (Figures 8B and 8C). End-diastolic LV internal dimensions, end-systolic LV internal dimensions, and LV mass showed no significant difference between saline- and E6446-treated TAC-operated mice until 3 weeks after TAC. However, the parameters were significantly smaller in E6446-treated mice than those in saline-treated mice 4 and 6 weeks after the operation. Fractional shortening was significantly higher in E6446-treated mice than that in saline-treated mice 6 weeks after TAC. There were no significant differences in IVSd and end-diastolic LV posterior wall thickness between the 2 groups at any time point. The heart weight-to-tibia length and lung weight-to-tibia length ratios were significantly reduced by E6446 treatment (Figure 8D). Schematic protocols to examine the effect of treatment of E6446 on cardiac phenotypes are described in Figures 9 and 10.

## DISCUSSION

The present study showed that E6446 prevents the development of pressure overload-induced heart failure when administered before the cardiac event and also suppresses the progression of heart failure when started after cardiac dysfunction manifested. We have reported that TLR9 is essential in producing inflammatory cytokines in failing hearts (7). TLRs are essential in driving the recruitment of inflammatory cells and production of cytokines during cardiac remodeling (13). E6446 prevents cellular events activated by TLR9, exerting broader inhibitory effects on inflammatory cytokine production, and thus the treatment of heart failure with the inhibitor has an advantage over the therapy neutralizing only 1 cytokine. The near-complete rescue of TAC-induced LV dilatation and dysfunction by E6446 pretreatment suggests that the TLR9-signaling pathway is the



dominant pathway for inducing adverse ventricular remodeling, with a limited role for other pathways such as nucleotide-binding domain leucine-rich repeat containing protein 3 (NLRP3) and cyclic GMP-AMP synthase (cGAS)-stimulator of interferon genes (STING) activation by mitochondrial DNA released in the cytosol in the setting of mitophagy dysfunction (14).

In human peripheral blood mononuclear cells or mouse spleen cells, E6446 diminished IL-6 production in response to CpG ODN (8). A 100-fold higher concentration of E6446 inhibited cytokine production in response to the imidazoquinoline compound R848, which is a TLR7/8 agonist (15). When C57BL/6 mice were orally treated with E6446, E6446 completely inhibited CpG ODN-induced IL-6 production in sera but not R848- and LPS-mediated cytokine production (8). Consistent with these results, our findings indicate that E6446 specifically

inhibited the expression of inflammatory cytokines through a TLR9-dependent pathway but not TLR4- or TLR7-dependent pathways in adult cardiomyocytes. Thus, E6446 has high specificity to TLR9. In mouse bone marrow-derived dendritic cells, E6446 potentially inhibited IL-6 production induced by CpG ODN but not by TLR3 ligands (9). However, E6446 was a potent inhibitor of IL-6 induction by single-stranded RNA, a TLR7/8 agonist, but a relatively poor inhibitor of IL-6 induction by R848, suggesting that the ability of E6446 to suppress TLR7/8 might be ligand dependent. Based on our data showing the importance of TLR9 signaling in the development of inflammation and heart failure and its specificity to TLR9 in cardiomyocytes, the cardioprotective action of E6446 is TLR9 mediated. However, we cannot exclude the possibility that TLR7/8 is involved in the effect of E6446 on the development of heart failure.

E6446 inhibits *in vitro* DNA–TLR9 interaction via an association with DNA but not with TLR9 (9). Furthermore, E6446 accumulates in the intracellular acidic compartment. Mitochondrial DNA is accumulated in autolysosome and coexists with TLR9 in failing hearts (7). DNase II activity was up-regulated in hypertrophied hearts but not in failing hearts. The incomplete digestion of mitochondrial DNA would be due to the loss of up-regulation of DNase II activity. Mitophagy impairment occurs in the TAC-induced mouse heart failure model (16). Thus, it is also possible that impairment of mitophagy or lysosomal permeabilization or lysosomal dysfunction might result in the accumulation of mitochondrial DNA in autolysosome. Our data in this study showed that there was no significant difference in the number of EdU and LAMP2a double-positive deposits between TAC-operated saline- and E6446-treated hearts. This outcome suggests that E6446 has no effect on mitochondrial DNA accumulation in autolysosomes. Thus, we can assume that the E6446 administered accumulates in lysosomes in the cardiomyocytes and interacts with mitochondrial DNA. When E6446 was orally administered to mice before TAC, E6446 inhibited TLR9 signaling by interfering with the mitochondrial DNA–TLR9 interaction and subsequent development of inflammation and heart failure.

We showed that *Tlr9*<sup>−/−</sup> mice are more resistant to pressure overload than control mice, and inhibitory ODN to TLR9 (ODN2088) improved survival in TAC-operated wild-type mice when administered before TAC (7). However, administration of a drug before cardiac events is not clinically relevant. The results indicate that E6446 can slow the development of heart failure even after cardiac dysfunction

manifested. Thus, E6446 or other immunomodulatory therapy can be used to prevent or delay pressure overload-induced heart failure.

**STUDY LIMITATIONS.** This study shows the therapeutic effects of a TLR9 inhibitor, E6446, on mouse pressure-overload heart failure model. Obviously, further studies are necessary to translate the findings into human heart failure therapy. TAC-induced mouse model does not fully represent the complex features of clinical heart failure. To establish the clinical feasibility of E6446 treatment for heart failure, the effects of E6446 on different heart failure models have to be examined, such as myocardial infarction. In addition, we used young healthy mice in this study. However, in most patients, and particularly in elderly patients, heart failure is accompanied by a range of comorbidities, such as hypertension, diabetes mellitus, renal dysfunction and hyperlipidemia. Such factors may influence on the cardioprotective effect of E6446 in heart failure. Thus, further investigation using various disease models is required to clarify the clinical target of E6446 treatment. Furthermore, it will be important to validate the findings in large animal models and ultimately in human patients. It remains unclear whether mitochondrial DNA is accumulated in autolysosomes and mitochondrial DNA–TLR9 axis is involved in the genesis of inflammation in human failing hearts.

## CONCLUSIONS

Heart failure is the result of various cardiac diseases such as myocardial infarction, high blood pressure, cardiomyopathy, valvular diseases, arrhythmia, and congenital heart diseases. Elevated levels of inflammatory mediators have been identified in patients with heart failure, including heart failure with reduced and preserved ejection fraction, as well as short-term decompensated heart failure (1). Thus, investigation of the involvement of the TLR9-signaling pathway in other mouse or larger animal heart failure models and various types of human heart failure is warranted. We ultimately will be able to identify subsets of patients with heart failure who will benefit from inhibition of TLR9 signaling.

**ACKNOWLEDGMENTS** The authors thank Eisai Inc. for the generous gift of E6446, and Ms. Yuri Kiura for technical assistance.

**ADDRESS FOR CORRESPONDENCE:** Dr. Kinya Otsu, School of Cardiovascular Medicine and Sciences, King's College London British Heart Foundation Centre of Excellence, 125 Coldharbour Lane, London SE5 9NU, United Kingdom. E-mail: [kinya.otsu@kcl.ac.uk](mailto:kinya.otsu@kcl.ac.uk).

## PERSPECTIVES

**COMPETENCY IN MEDICAL KNOWLEDGE:** Heart failure is a major health threat in the developed countries with high morbidity and mortality. Novel and effective therapeutic agents against heart failure need to be developed. Inflammation and proinflammatory cytokines play an important role in the pathogenesis of heart failure. Inflammatory mediators can be therapeutic targets in heart failure.

**TRANSLATIONAL OUTLOOK:** A TLR9 inhibitor, E6446, exerted a beneficial effect on attenuating the development or progression of heart failure in a pressure overload-induced mouse model. E6446 treatment has an advantage over targeted anticytokine approaches using biological response modifiers, because it modulates a broad spectrum of inflammatory mediators. Thus, it may be a new promising therapeutic agent for human heart failure.

## REFERENCES

1. Mann DL. Innate immunity and the failing heart: the cytokine hypothesis revisited. *Circ Res* 2015;116:1254–68.
2. Chung ES, Packer M, Lo KH, Fasanmade AA, Willerson JT. Randomized, double-blind, placebo-controlled, pilot trial of infliximab, a chimeric monoclonal antibody to tumor necrosis factor- $\alpha$ , in patients with moderate-to-severe heart failure: results of the anti-TNF Therapy Against Congestive Heart Failure (ATTACH) trial. *Circulation* 2003;107:3133–40.
3. Mann DL, McMurray JJ, Packer M, et al. Targeted anticytokine therapy in patients with chronic heart failure: results of the Randomized Etanercept Worldwide Evaluation (RENEWAL). *Circulation* 2004;109:1594–602.
4. Deng GM, Nilsson IM, Verdrengh M, Collins LV, Tarkowski A. Intra-articularly localized bacterial DNA containing CpG motifs induces arthritis. *Nat Med* 1999;5:702–5.
5. Hemmi H, Takeuchi O, Kawai T, et al. A Toll-like receptor recognizes bacterial DNA. *Nature* 2000;408:740–5.
6. Saito T, Sadoshima J. The molecular mechanisms of mitochondrial autophagy/mitophagy in the heart. *Circ Res* 2015;116:1477–90.
7. Oka T, Hikoso S, Yamaguchi O, et al. Mitochondrial DNA that escapes from autophagy causes inflammation and heart failure. *Nature* 2012;485:251–5.
8. Franklin BS, Ishizaka ST, Lamphier M, et al. Therapeutic targeting of nucleic acid-sensing Toll-like receptors prevents experimental cerebral malaria. *Proc Natl Acad Sci U S A* 2011;108:3689–94.
9. Lamphier M, Zheng W, Latz E, et al. Novel small molecule inhibitors of TLR7 and TLR9: mechanism of action and efficacy in vivo. *Mol Pharmacol* 2014;85:429–40.
10. Nakai A, Yamaguchi O, Takeda T, et al. The role of autophagy in cardiomyocytes in the basal state and in response to hemodynamic stress. *Nat Med* 2007;13:619–24.
11. Hoshino K, Takeuchi O, Kawai T, et al. Cutting edge: Toll-like receptor 4 (TLR4)-deficient mice are hyporesponsive to lipopolysaccharide: evidence for TLR4 as the *lps* gene product. *J Immunol* 1999;162:3749–52.
12. Florian H, Parviz AN, Hiroaki H, et al. The Toll-like receptor 7 (TLR7)-specific stimulus loxoribine uncovers a strong relationship within the TLR7, 8 and 9 subfamily. *Eur J Immunol* 2003;33:2987–97.
13. Mann DL. Inflammatory mediators and the failing heart: past, present, and the foreseeable future. *Circ Res* 2002;91:988–98.
14. Nakayama H, Otsu K. Mitochondrial DNA as an inflammatory mediator in cardiovascular diseases. *Biochem J* 2018;475:839–52.
15. Jurk M, Heil F, Vollmer J, et al. Human TLR7 or TLR8 independently confer responsiveness to the antiviral compound R-848. *Nat Immunol* 2002;3:499.
16. Shirakabe A, Zhai P, Ikeda Y, et al. Drp1-dependent mitochondrial autophagy plays a protective role against pressure overload-induced mitochondrial dysfunction and heart failure. *Circulation* 2016;133:1249–63.

**KEY WORDS** heart failure, mitochondria, pressure overload, Toll-like receptor 9